

Extraction of the strong coupling constant and strange quark mass from semileptonic τ decays¹

A. A. Pivovarov

Institut für Physik der Johannes-Gutenberg-Universität,
Staudinger Weg 7, 55099 Mainz, Germany
and

Institute for Nuclear Research of the Russian Academy of Science,
117312 Moscow, Russia

Abstract

In this lecture I present a pedagogical introduction to the low-energy phenomenology of light flavors. The renormalization scheme freedom in defining QCD parameters is discussed. It is shown in some details how one can extract an accurate numerical value for the strong coupling constant from the τ -lepton decay rate into hadrons. As a related topic I discuss some peculiarities of definition of the quark mass in theories with confinement and describe the strange quark mass determination from data on τ -lepton decays employing contour resummation which is a modern technique of the precision analysis in perturbative QCD.

1 Definition of QCD parameters α_s and m_s

Quantum ChromoDynamics (QCD) as a field theory for describing strong interactions is given by the Lagrangian

$$\mathcal{L}_{\text{QCD}} = \sum_f \bar{q}_f (i\gamma_\mu \partial^\mu + g_s \gamma^\mu G_\mu^a t^a - m_f) q_f - \frac{1}{4} G_{\mu\nu}^a G_{\mu\nu}^a$$

where G_μ^a is a non-Abelian gluonic field and $G_{\mu\nu}^a$ is the field strength tensor [1]. There are six quark flavors $q = (u, d, s, c, b, t)$, three of which (u, d, s) are called light while the other three are heavy. There is a close analogy with QED – the Abelian gauge theory for describing the electromagnetic interaction of charged leptons. The QED Lagrangian for charged leptons reads

$$\mathcal{L}_{\text{QED}} = \sum_l \bar{l} (i\gamma_\mu \partial^\mu + e\gamma^\mu A_\mu - m_l) l - \frac{1}{4} F_{\mu\nu} F_{\mu\nu}$$

where $F_{\mu\nu}$ is the electromagnetic field strength tensor. In the standard model there are three charged leptons $l = (e, \mu, \tau)$: electron and muon are very light at the hadronic mass scale of order 1 GeV while the τ lepton is rather heavy with a mass $M_\tau = 1.777$ GeV [2].

The interaction is given by a vertex in the Lagrangian and normalized to a coupling constant g_s at a tree level of perturbation theory (PT). A full theory (beyond the tree level) introduces

¹Lecture given at International School “Heavy Quark Physics” Dubna, May 28 - June 5, 2002

a dressed vertex that eventually determines a coupling constant after renormalization. For the renormalizable models of quantum field theory as QED and QCD the PT dressing is straightforward and can in principle be done at any finite order of the expansion in the coupling constant. However, the dressing procedure is not unique because renormalization of loops can introduce some freedom in the choice of finite parameters through a particular definition of counterterms [3].

Leptons can be detected as asymptotic states in the scattering processes that allows one to relate the coupling – for instance, the fine structure constant α – and the particle mass m_l to observables very directly. Thus, the electron-photon scattering at low energies can be used to define the coupling constant (the fine structure constant, in fact) $\alpha = e^2/4\pi$ through the Thompson cross section. This is a natural definition directly through a physical observable. Lepton mass is also directly related to experiment: it can be defined as a physical mass of the asymptotic state (a position of the pole of the lepton propagator). In QCD there is a phenomenon of confinement and no asymptotic states of quarks and/or gluons can be observed. Only the colorless hadrons appear as the asymptotic states. Therefore, definitions of the coupling constant and quark masses in QCD are less direct than in QED. At the theoretical level of the given Lagrangian they are very similar though: a vertex for the coupling and the propagator for a mass. To determine numerical values for the coupling and quark masses one should turn to experiment. As there is no possibility to measure these quantities directly one should specify the research area as the definitions of the parameters can be adopted to specific experiments (basically to a corresponding energy scale). In this lecture I will talk about τ -lepton physics which is the area of low-energy hadron phenomenology: hadronic states have an energy $E < M_\tau$. The particles which can be observed, for instance, in the process of e^+e^- annihilation are ρ , ω , φ , $\pi\pi$. In τ decays $\tau \rightarrow \nu + \text{hadrons}$ one can in addition see π , $a_1(1260)$, K , $K^*(892)$.

The primary difficulty for extracting the QCD coupling from experimental data is that the $q\bar{q}g$ vertex cannot be “directly” measured. Indeed, a typical process (vertex) with hadrons at low energies is $\rho \rightarrow \pi\pi$ which is not directly expressed theoretically through the quark-gluon interaction vertex. Therefore, extracting α_s (defined in terms of quark-gluon vertex in the Lagrangian) from the experimental quantity as, for instance, the ρ -meson decay width $\Gamma(\rho \rightarrow \pi\pi)$ is highly nontrivial. One can also see a propagation of the pion but not that of a quark. No mass shell for the quark (or gluon) is seen in the experiment. Thus, in QCD there is no preferable definition of parameters related to experiment. Then the only guidance for the choice of a particular definition of the QCD parameters in PT is the technical convenience (and also some general requirements such as gauge invariance) [4].

Presently dimensional regularization is overwhelmingly used in many loop calculations. The renormalization procedure is usually a minimal one – subtraction of poles in $\varepsilon = (D-4)/2$ where D is space-time dimension (an arbitrary complex number formally introduced for the regularization purposes). This procedure is quite abstract and remote from experimental quantities. Thus, formally, the $\overline{\text{MS}}$ -scheme coupling constant is defined by $\alpha_s^{\overline{\text{MS}}}(\mu) = Z_\alpha^{\overline{\text{MS}}}(\mu)\alpha_s^B$ where α_s^B is a bare coupling constant. This definition of the renormalized coupling constant is not unique. In the momentum subtraction (MOM) scheme $\alpha_s^{\text{MOM}}(\mu) \sim \Gamma_{\text{qdg}}^{\text{ren}}(p_1^2 = p_2^2 = p_3^2 = \mu^2)$. Since the definition is not unique the renormalization scheme freedom emerges. It is controlled by the renormalization group and can be conveniently parameterized by the coefficients $\beta_{2,3,\dots}$ of the β -function [5, 6]

$$\mu \frac{\partial}{\partial \mu} \alpha_s(\mu) = 2\beta(\alpha_s), \quad \beta(\alpha_s) = -\beta_0 \alpha_s^2 - \beta_1 \alpha_s^3 - \beta_2 \alpha_s^4 - \beta_3 \alpha_s^5 + O(\alpha_s^6)$$

and, therefore, any particularly defined coupling constant α_s^{sch} generates an associated $\beta^{\text{sch}}(\alpha_s^{\text{sch}})$ -

function. The same is true for the definition of the quark mass. In the $\overline{\text{MS}}$ -scheme one defines the renormalized mass as $m^{\overline{\text{MS}}}(\mu) = Z_m^{\overline{\text{MS}}}(\mu)m^B$ with an associated γ -function $(\mu\partial/\partial\mu)m(\mu) = 2\gamma(\alpha_s)m(\mu)$. Functions $\beta(\alpha_s)$ and $\gamma(\alpha_s)$ are known up to four-loop approximation in PT [7]. The pole mass is difficult to define for light quarks since numerically even the strange quark is very light $m_s \sim \Lambda_{\text{QCD}}$.

In fact, a QCD coupling constant can also be defined beyond PT through physical observables. For light flavor phenomenology it can directly be defined through the cross section of e^+e^- annihilation

$$1 + \alpha_s(s) \sim \frac{\sigma(e^+e^- \rightarrow \text{hadrons})}{\sigma(e^+e^- \rightarrow \mu\bar{\mu})}$$

while for heavy quark physics the definition based on the heavy-quark static potential $\alpha_V(\bar{q}^2) \sim V_{q\bar{q}}(\bar{q}^2)$ can be useful [8].

2 Kinematics of semileptonic τ decays

The differential decay rate of the τ lepton into an hadronic system $H(s)$ with a total squared energy s

$$\frac{d\sigma(\tau \rightarrow \nu H(s))}{ds} \sim \left(1 - \frac{s}{M_\tau^2}\right)^2 \left(1 + \frac{2s}{M_\tau^2}\right) \rho(s)$$

is determined by the hadronic spectral density $\rho(s)$ defined through the correlator of weak currents. For the (ud) current $j_\mu^W(x) = \bar{u}\gamma_\mu(1 - \gamma_5)d$ one finds

$$i \int \langle T j_\mu^W(x) j_\nu^{W+}(0) \rangle e^{iqx} dx = (q_\mu q_\nu - q^2 g_{\mu\nu}) \Pi^{\text{had}}(q^2), \quad \Pi^{\text{had}}(q^2) = \int \frac{\rho(s) ds}{s - q^2} \quad (1)$$

with $\rho(s) \sim \text{Im } \Pi^{\text{had}}(s + i0)$, $s = q^2$. The function $\Pi^{\text{had}}(Q^2)$ with $Q^2 = -q^2$ is calculable in pQCD far from the physical cut as a series in the running coupling constant $\alpha_s(Q^2)$.

NonPT effects (power corrections) are included using OPE for the correlator (1) at small distances as $x \rightarrow 0$ in Euclidean domain (that corresponds to large Q^2) through phenomenological characteristics of the vacuum such as gluon and quark condensates [9]. The lattice approximation for the evaluation of the correlator $\Pi^{\text{had}}(Q^2)$ beyond PT can also be used [10]. This is a basis for theoretical description of semileptonic τ decays in QCD.

Integrating the function $\Pi^{\text{had}}(z)$ over a contour in the complex q^2 plane beyond the physical cut $s > 0$ one finds that for particular weight functions some integrals of the hadronic spectral density $\rho(s)$ can be reliably computed in PT [11, 12]. Indeed, due to Cauchy theorem one gets

$$\oint_C \Pi(z) dz = \int_{\text{cut}} \rho(s) ds.$$

Using the approximation $\Pi^{\text{had}}(z)|_{z \in C} \approx \Pi^{\text{PT}}(z)|_{z \in C}$ which is well justified sufficiently far from the physical cut one obtains

$$\oint_C \Pi^{\text{had}}(z) dz = \int_{\text{cut}} \rho(s) ds = \oint_C \Pi^{\text{PT}}(z) dz$$

i.e. the integral over the hadronic spectrum can be evaluated in pQCD. The total decay rate of the τ lepton written in the form of an integral along the cut

$$R_{\tau S=0} = \frac{\Gamma(\tau \rightarrow H_{S=0}\nu)}{\Gamma(\tau \rightarrow l\bar{\nu}\nu)} \sim \int_{\text{cut}} \left(1 - \frac{s}{M_\tau^2}\right)^2 \left(1 + \frac{2s}{M_\tau^2}\right) \rho(s) ds$$

is precisely the quantity that one can reliably compute in pQCD [13].

3 PT analysis in QCD

For technical reasons (no overall UV divergence) a derivative of the correlator $\Pi^{\text{had}}(Q^2)$ is often used for presenting results of PT evaluation

$$D(Q^2) = -Q^2 \frac{d}{dQ^2} \Pi^{\text{had}}(Q^2) = Q^2 \int \frac{\rho(s) ds}{(s + Q^2)^2}.$$

The PT expansion for the D -function in terms of $a_s(Q) = \alpha_s(Q)/\pi$ reads $D(Q^2) = 1 + a_s(Q) + k_1 a_s(Q)^2 + k_2 a_s(Q)^3 + k_3 a_s(Q)^4$. In the $\overline{\text{MS}}$ -scheme

$$k_1 = \frac{299}{24} - 9\zeta(3), \quad k_2 = \frac{58057}{288} - \frac{779}{4}\zeta(3) + \frac{75}{2}\zeta(5)$$

with ζ -function equal $\zeta(3) = 1.202\dots$ and $\zeta(5) = 1.037\dots$. The above numerical values for $k_{1,2}$ summarize the results of three and four loop PT calculations [14]. Numerically, one finds $D(Q^2) = 1 + a_s + 1.64a_s^2 + 6.37a_s^3 + k_3a_s^4$. Coefficient k_3 is known only partly [15]. It is retained to obtain a feeling for the possible magnitude of the $O(\alpha_s^4)$ correction. The decay rate of the τ lepton into nonstrange hadrons is written in the form

$$R_{\tau S=0} = \frac{\Gamma(\tau \rightarrow H_{S=0}\nu)}{\Gamma(\tau \rightarrow l\bar{\nu}\nu)} = 3|V_{ud}|^2(1 + \delta_P + \delta_{NP}).$$

Here the first term is the parton model result, the second term δ_P represents pQCD effects. NonPT effects are small, $\delta_{NP} \approx 0$, in the factorization approximation for the four-quark vacuum condensates which is quite accurate [13, 16].

The experimental result $R_{\tau S=0}^{\text{exp}} = 3.492 \pm 0.016$ leads to $\delta_P^{\text{exp}} = 0.203 \pm 0.007$ [17]. In the $\overline{\text{MS}}$ -scheme the correction δ_P is given by the series

$$\delta_P^{\text{th}} = a_s + 5.2023a_s^2 + 26.366a_s^3 + (78.003 + k_3)a_s^4 + O(a_s^5)$$

with a_s taken at the scale $\mu = M_\tau$. Usually one extracts a numerical value for $\alpha_s(M_\tau)$ by treating the first three terms of the expression as an exact function – the cubic polynomial $a_s + 5.2023a_s^2 + 26.366a_s^3 = \delta_P^{\text{exp}}$. The solution reads $\pi a_s^{\text{st}}(M_\tau) \equiv \alpha_s^{\text{st}}(M_\tau) = 0.3404 \pm 0.0073_{\text{exp}}$. The error is due to the error of the input experimental value δ_P^{exp} . It is difficult to estimate the theoretical uncertainty of the approximation for the (asymptotic) series given by the cubic polynomial (higher order terms). One criterion is the pattern of convergence of the series

$$\delta_P^{\text{exp}} = 0.203 = 0.108 + 0.061 + 0.034 + \dots$$

The corrections provide a 100% change of the leading term. Another criterion is the order-by-order behavior of the extracted numerical value for the coupling constant. In consecutive orders of PT

$$\alpha_s^{\text{st}}(M_\tau)_{LO} = 0.6377, \quad \alpha_s^{\text{st}}(M_\tau)_{NLO} = 0.3882, \quad \alpha_s^{\text{st}}(M_\tau)_{NNLO} = 0.3404$$

that translates into a series for the coupling constant

$$\alpha_s^{\text{st}}(M_\tau)_{NNLO} = 0.6377 - 0.2495 - 0.0478 - \dots$$

One can take a half(?) of the last term as an estimate of the theoretical uncertainty. It is only an indicative estimate. No rigorous justification can be given for such an assumption about the accuracy of the approximation without knowledge of the structure of the whole series. The

uncertainty obtained in such a way $\Delta\alpha_s^{st}(M_\tau)_{th} = 0.0478/2 = 0.0239 \gg 0.0073_{exp}$ is much larger than that from experiment. This is a challenge for the theory: the accuracy of theoretical formulae cannot compete with experimental precision. Assuming this theoretical uncertainty one has

$$\alpha_s^{st}(M_\tau)_{NNLO} = 0.3404 \pm 0.0239_{th} \pm 0.0073_{exp}$$

Theory dominates the error. Still it is not the whole story. Now one can choose a different expansion parameter. The simplest way is to change the scale of the coupling along the RG trajectory $M_\tau \rightarrow 1 \text{ GeV}$. In terms of $a_s(1 \text{ GeV})$ one finds a series $a_s(1) + 2.615a_s(1)^2 + 1.54a_s(1)^3 = \delta_P^{\text{exp}}$. The solution for the coupling constant is $\alpha_s(1 \text{ GeV}) = 0.453$. The convergence pattern for the correction δ_P^{exp} is

$$\delta_P^{\text{exp}} = 0.203 = 0.144 + 0.054 + 0.005$$

and for the numerical value of the coupling constant

$$\alpha_s(1) = 0.453 = 0.638 - 0.177 - 0.008.$$

Should one conclude that now the accuracy is much better? What would be an invariant criterion for the precision of theoretical predictions obtained from PT, i.e. finite number of terms of asymptotic series? Thus, one sees that the renormalization scheme dependence can strongly obscure the heuristic evaluation of the accuracy of theoretical formulae in the absence of any information on the structure of the whole PT series. The final result for the standard reference value of the coupling constant normalized to the Z boson mass M_Z reads [18]

$$\alpha_s(M_Z)_\tau = 0.1184 \pm 0.0007_{exp} \pm 0.0006_{hq \text{ mass}} \pm 0.0010_{th=truncation}$$

with a theoretical uncertainty that mainly comes from the truncation of the PT series. The world average value given by the Particle Data Group reads [2]

$$\alpha_s(M_Z)_{\text{PDG}}^{\text{av}} = 0.1172 \pm 0.002.$$

To reduce a renormalization scheme dependence of the theoretical analysis one should use several observables simultaneously [19]. Such a possibility has recently emerged in study of τ decays since experimental data on Cabibbo suppressed ($S = 1$) channel appeared [20]. For strange hadrons $H_{S=1}$ (us part of the weak current: Cabibbo suppressed decays) the decay rate becomes

$$R_{\tau S=1} = \frac{\Gamma(\tau \rightarrow H_{S=1}\nu)}{\Gamma(\tau \rightarrow l\bar{\nu}\nu)} = 3|V_{us}|^2(1 + \delta'_P + \delta'_{NP}).$$

The first term (“1”) is the parton model result, the second term δ'_P gives pQCD effects. Small s -quark mass effects for Cabibbo suppressed part of the rate are taken in PT at the leading order in the ratio m_s^2/M_τ^2 since $m_s \ll M_\tau$

$$\delta'_P(\alpha_s, m_s) = \delta_P(\alpha_s) + \frac{m_s^2}{M_\tau^2} \Delta_m(\alpha_s)$$

with $\delta_P(\alpha_s)$ being a correction in massless approximation for light quarks that is well justified for nonstrange decays (ud part) since u, d quarks are very light indeed $m_u + m_d = 14 \text{ MeV}$ [21, 22]. The correlator of the weak charged strange current $j_\mu(x) = \bar{u}\gamma_\mu(1 - \gamma_5)s$ with a finite s -quark mass is not transverse

$$i \int dx e^{iqx} \langle T j_\mu(x) j_\nu^\dagger(0) \rangle = q_\mu q_\nu \Pi_q(q^2) + g_{\mu\nu} \Pi_g(q^2).$$

Retaining the first order term of expansion in the small ratio m_s^2/q^2 one finds the m_s^2 correction to the invariant functions $\Pi_{q,g}(q^2)$

$$\Pi_q(q^2) = \Pi(q^2) + 3\frac{m_s^2}{q^2}\Pi_{mq}(q^2), \quad \Pi_g(q^2) = -q^2\Pi(q^2) + \frac{3}{2}m_s^2\Pi_{mg}(q^2)$$

where $\Pi(q^2)$ is an invariant function for the mass zero case. The functions $\Pi_{q,g}(Q^2)$ are computable in QCD perturbation theory within operator product expansion for $Q^2 \rightarrow \infty$. Thus, the experimental data on τ lepton decays are theoretically described by three independent invariant functions (form factors) which can be analyzed simultaneously that may help to reduce uncertainties introduced by the renormalization scheme freedom.

In the actual analysis one can factor out the renormalization scheme freedom to large extent by introducing an effective scheme with definitions of effective quantities a , m_q^2 , m_g^2 through the relations [23]

$$\begin{aligned} -Q^2 \frac{d}{dQ^2} \Pi(Q^2) &= 1 + a(Q^2), \\ -m_s^2(M_\tau^2) Q^2 \frac{d}{dQ^2} \Pi_{mg}(Q^2) &= m_g^2(M_\tau^2) C_g(Q^2), \\ m_s^2(M_\tau^2) \Pi_{mq}(Q^2) &= m_q^2(M_\tau^2) C_q(Q^2). \end{aligned} \quad (2)$$

Here $C_{q,g}(Q^2)$ are coefficient functions of mass corrections. They are conveniently normalized by the requirement $C_{q,g}(M_\tau^2) = 1$. In terms of the $\overline{\text{MS}}$ scheme quantities $\alpha_s \equiv \alpha_s(M_\tau^2)$ and $m_s \equiv m_s(M_\tau^2)$ the effective parameters in eq. (2) read

$$\begin{aligned} a(M_\tau^2) &= \frac{\alpha_s}{\pi} + k_1 \left(\frac{\alpha_s}{\pi} \right)^2 + k_2 \left(\frac{\alpha_s}{\pi} \right)^3 + k_3 \left(\frac{\alpha_s}{\pi} \right)^4 + \mathcal{O}(\alpha_s^5), \\ m_g^2(M_\tau^2) &= m_s^2(M_\tau^2) \left(1 + \frac{5}{3} \frac{\alpha_s}{\pi} + k_{g1} \left(\frac{\alpha_s}{\pi} \right)^2 + k_{g2} \left(\frac{\alpha_s}{\pi} \right)^3 + \mathcal{O}(\alpha_s^4) \right), \\ m_q^2(M_\tau^2) &= m_s^2(M_\tau^2) \left(1 + \frac{7}{3} \frac{\alpha_s}{\pi} + k_{q1} \left(\frac{\alpha_s}{\pi} \right)^2 + k_{q2} \left(\frac{\alpha_s}{\pi} \right)^3 + \mathcal{O}(\alpha_s^4) \right). \end{aligned}$$

Numerical values for the coefficients k_3 , k_{q2} are unknown though their estimates within various intuitive approaches can be found in the literature [24].

For confronting the experimental data obtained in τ decays with theory of strong interactions one uses special integrals of the hadronic spectral density $\rho(s)$ (called spectral moments) of the form

$$\mathcal{M}_{kl}^{\text{exp}} = \int_0^{M_\tau^2} \rho(s) \left(1 - \frac{s}{M_\tau^2} \right)^k \left(\frac{s}{M_\tau^2} \right)^l \frac{ds}{M_\tau^2}$$

which is suitable for experiment as the spectrum $\rho(s)$ is measured. Another representation of the moments is suitable for the perturbation theory evaluation in QCD. Due to analytic properties of the functions $\Pi_{q,g}(Q^2)$ the moments can be rewritten as contour integrals in the complex q^2 plane:

$$\mathcal{M}_{kl}^{\text{th}} = \frac{i}{2\pi} \oint_C \Pi_{\#}(z) \left(1 - \frac{z}{M_\tau^2} \right)^k \left(\frac{z}{M_\tau^2} \right)^l \frac{dz}{M_\tau^2}.$$

Technically one chooses a circular contour in the complex Q^2 -plane with $Q^2 = M_\tau^2 e^{i\phi}$, $-\pi < \phi < \pi$ that converts all invariant amplitudes $\Pi_{q,g}(Q^2)$ to the certain functions of the angle ϕ . The evolution of the functions $\Pi_{q,g}(Q^2)$ along the contour in the complex plane is governed by the renormalization group [25]. For the massless case corresponding to the analysis of data in Cabibbo




Figure 1: Running of the functions $a(\phi)$ and $C_q(\phi)$ on a circular contour in the complex plane calculated at LO, NLO and NNLO (left: $a(\phi)$; right: $C_q(\phi)$; real parts only)

favorable channel the only relevant quantity is the running “coupling constant” $a(Q^2) \rightarrow a(\phi)$ for $Q^2 = M_\tau^2 e^{i\phi}$ that serves as the expansion parameter for the function $\Pi(Q^2)$ along the contour. Note that there are no higher order corrections in the effective scheme by definition. The renormalization group evolution is determined by the effective β function for the effective coupling constant [26, 27]

$$-i \frac{d}{d\phi} a(\phi) = \beta(a(\phi)), \quad a(\phi = 0) = a(M_\tau^2),$$

and the anomalous dimension for the running mass

$$-i \frac{d}{d\phi} m_s(\phi) = \gamma_m[a(\phi)] m_s(\phi), \quad m_s(\phi = 0) = m_s.$$

The initial values for $a(\phi)$ and $m_s(\phi)$ are extracted from fit to data. Thus, we have three quantities $a(Q)$, $C_q(Q)$, $C_g(Q)$ associated with three invariant functions $\Pi(Q^2)$, $\Pi_{q,g}(Q^2)$ describing the τ system in the considered approximation.

The RG equations for the set of quantities $\{a(Q), C_q(Q), C_g(Q)\}$ are

$$Q^2 \frac{d}{dQ^2} a(Q^2) = \beta(a), \quad Q^2 \frac{d}{dQ^2} C_{g,q}(Q^2) = 2\gamma_{g,q}(a) C_{g,q}(Q^2).$$

The RG functions $\beta(a)$ and $\gamma_{g,q}(a)$ are given by the expressions [23]

$$\frac{-4\beta(a)}{9a^2} = 1 + 1.778a + 5.24a^2 + a^3(-34 + 2k_3);$$

$$\frac{-\gamma_g(a)}{a} = 1 + 4.03a + 17.45a^2 + a^3(249.59 - k_3);$$

$$\frac{-\gamma_q(a)}{a} = 1 + 4.78a + 32.99a^2 + a^3(-252 - k_3 + 3.4k_{q2}).$$

The solution of the renormalization group equation for the effective coupling constant $a(\phi)$ converges well when the higher order corrections of the β -function are included. The change from the next-to-leading order (NLO) solution to the next-to-next-to-leading order (NNLO) is small. The behavior of the coefficient function $C_g(\phi)$ (not shown) related to the contributions of spin one particles is rather similar to that of the coupling constant. However, the convergence pattern of the function $C_q(\phi)$ is much worse (see Fig. 1). It seems that the γ_q -function has already shown

Table 1: Coefficients of eq. (3)

(k, l)	A_{kl}^{LO}	A_{kl}^{NLO}	A_{kl}^{NNLO}	B_{kl}^{LO}	B_{kl}^{NLO}	B_{kl}^{NNLO}
$(0, 0)$	1.361	1.445	1.434	0.523	0.601	0.625
$(1, 0)$	1.568	1.843	1.976	0.441	0.552	0.601
$(2, 0)$	1.762	2.282	2.646	0.390	0.530	0.607

up an asymptotic growth in the next-to-next-to-leading order which will limit the precision of our results. Note that the fact that the function $C_q(\phi)$ can behave wilder in higher order of perturbation theory is expected since this function is more infrared sensitive than the coupling constant $a(\phi)$ and the function $C_g(\phi)$.

The numerical value for the s -quark mass m_s is extracted from the difference between moments of Cabibbo-favored (ud -type) and Cabibbo-suppressed (us -type) decay rates

$$\delta R_\tau^{kl} = \frac{R_{\tau S=0}^{kl}}{|V_{ud}|^2} - \frac{R_{\tau S=1}^{kl}}{|V_{us}|^2}, \quad R_{\tau S=0,1}^{kl} = \int_0^{M_\tau^2} ds \left(1 - \frac{s}{M_\tau^2}\right)^k \left(\frac{s}{M_\tau^2}\right)^l \frac{dR_{\tau S=0,1}}{ds}.$$

The theoretical expression for the m_s^2 corrections to the moments (k, l) corresponding to the above experimental quantity is given by the contour integral in the complex q^2 plane

$$\frac{3im_s^2}{\pi} \oint_C \left(1 - \frac{z}{M_\tau^2}\right)^{2+k} \left(\frac{z}{M_\tau^2}\right)^l \left(\frac{\Pi_{mq}(z)}{z} - \frac{\Pi_{mg}(z)}{M_\tau^2}\right) \frac{dz}{M_\tau^2}.$$

In the theoretical expression for the difference δR_τ^{kl} we neglect terms of the order m_s^3/M_τ^3 , set the u - and d -quark masses to zero, and retain only the most important term linear in m_s . Within operator product expansion the coefficient of this term is given by the quark condensate. The final result for the difference reads [23]

$$\delta R_\tau^{kl} = 3S_{EW} \left(6 \frac{m_s^2}{M_\tau^2} (\omega_q A_{kl} + \omega_g B_{kl}) - 4\pi^2 \frac{m_s}{M_\tau} \frac{\langle \bar{s}s \rangle}{M_\tau^3} T_{kl} \right) \quad (3)$$

with $S_{EW} = 1.0194$ [28]. Here $m_{q,g}^2 = \omega_{q,g} m_s^2$ with $\omega_q = 1.73 \pm 0.04$, $\omega_g = 1.42 \pm 0.03$. We use the relation between vacuum condensates of strange and nonstrange quarks

$$\langle \bar{s}s \rangle = (0.8 \pm 0.2) \langle \bar{u}u \rangle$$

and the numerical value $\langle \bar{u}u \rangle = -(0.23 \text{ GeV})^3$ [29]. In the leading order approximation of the QCD perturbation theory for the coefficient function of the quark condensate the quantities T_{kl} multiplying the quark condensate are given by the expression

$$T_{kl} = 2(\delta_{l,0}(k+2) - \delta_{l,1}).$$

The numerical values for the first few coefficients T_{kl} read

$$T_{00} = 4, \quad T_{10} = 6, \quad T_{20} = 8, \quad T_{01} = -2, \quad T_{11} = -2.$$

The numerical values for the coefficients A_{kl} and B_{kl} are given in Table 1.

Table 2: Experimental moments and extracted mass

(k, l)	$(\delta R_\tau^{kl})^{\text{exp}}$	$m_s(M_\tau^2)$ MeV
(0, 0)	0.394 ± 0.137	$130 \pm \delta_{00}^{\text{th}} (= 6)$
(1, 0)	0.383 ± 0.078	$111 \pm \delta_{10}^{\text{th}} (= ?)$
(2, 0)	0.373 ± 0.054	$95 \pm \delta_{20}^{\text{th}} (= ?)$

Using experimental data one extracts m_s as given in Table 2. Theoretical prediction for the moment (0, 0) is the most reliable from PT point of view as $\delta_{20}^{\text{th}} (= ?) > \delta_{10}^{\text{th}} (= ?) > \delta_{00}^{\text{th}} = 6$ MeV. The final result reads

$$m_s(M_\tau^2) = 130 \pm 27_{\text{exp}} \pm 3_{\langle \bar{s}s \rangle} \pm 6_{\text{th}} \text{ MeV}.$$

Normalization at 1 GeV gives

$$m_s(1 \text{ GeV}) = 176 \pm 37_{\text{exp}} \pm 4_{\langle \bar{s}s \rangle} \pm 9_{\text{th}} \text{ MeV}.$$

Only the moment (0, 0) is used for the m_s determination as the most reliable one from the PT point of view. The higher order moments with the weight function $(1 - s/M_\tau^2)^k$ for large k have an uncontrollable admixture of higher dimension condensates that makes them strongly nonperturbative and, therefore, unreliable for applications based on PT calculations [30]. The contributions of higher dimension condensates are unknown and from general considerations the errors $\delta_{20}^{\text{th}} (= ?) > \delta_{10}^{\text{th}} (= ?)$ are expected to be much larger than $\delta_{00}^{\text{th}} = 6$ MeV. The value of the strange quark mass obtained by using the effective scheme approach as described in the present paper is in a reasonable agreement with other estimates [31, 32, 33]. It is a bit larger than the recent lattice determination [34].

To conclude, the experimental information on τ decays is a reliable source for the precision determination of the numerical values of important QCD parameters α_s and m_s .

Acknowledgements

I thank K.G. Chetyrkin, S. Groote, J.G. Körner, J.H. Kühn for useful discussions and fruitful collaboration. This work is partially supported by the Russian Fund for Basic Research under contracts 01-02-16171 and 02-01-00601 and by the Volkswagen grant.

References

- [1] C.N. Yang and R.L. Mills: Phys. Rev. **96**, 191 (1954); M. Gell-Mann: Phys. Lett. **8**, 214 (1964); M-Y. Han, Y. Nambu: Phys. Rev. B **139**, 1006 (1965); W.A. Bardeen, H. Fritzsch, M. Gell-Mann: In: *Scale and Conformal Symmetry in Hadron Physics*. ed. by R. Gatto (Wiley, New York 1973); H. Pagels: Phys. Rept. C **16**, 219 (1975)
- [2] D.E. Groom et al. (Particle Data Group): Eur. Phys. J. C **15**, 1 (2000)
- [3] N.N. Bogoliubov and D.V. Shirkov: *Quantum Fields* (Benjamin, New York 1983)
- [4] I.I. Bigi, M.A. Shifman, N.G. Uraltsev and A.I. Vainshtein: Phys. Rev. D **50**, 2234 (1994); M. Beneke and V.M. Braun: Nucl. Phys. B **426**, 301 (1994)

- [5] E. Stuekelberg and A. Peterman: *Helv. Phys. Acta* **26**, 499 (1953); M. Gell-Mann and E.E. Low: *Phys. Rev.* **95**, 1300 (1954); N.N. Bogoliubov and D.V. Shirkov: *Dokl. Akad. Nauk* **103**, 391 (1955)
- [6] P.M. Stevenson: *Phys. Rev. D* **23**, 2916 (1981)
- [7] T. van Ritbergen, J.A.M. Vermaseren and S.A. Larin: *Phys. Lett. B* **400**, 379 (1997); K.G. Chetyrkin: *Phys. Lett. B* **404**, 161 (1997); T. van Ritbergen, J.A.M. Vermaseren and S.A. Larin: *Phys. Lett. B* **405**, 327 (1997)
- [8] M. Jezabek, J.H. Kuhn, M. Peter, Y. Sumino and T. Teubner: *Phys. Rev. D* **58**, 014006 (1998); N.V. Krasnikov and A.A. Pivovarov: *Mod. Phys. Lett. A* **11**, 835 (1996); *Phys. Atom. Nucl.* **64**, 1500 (2001); M. Luscher, R. Sommer, P. Weisz and U. Wolff: *Nucl. Phys. B* **413**, 481 (1994); A.A. Pivovarov: “Heavy quark production near the threshold in QCD.” [arXiv:hep-ph/0110398]
- [9] H.D. Politzer: *Nucl. Phys. B* **117**, 397 (1976); M.A. Shifman, A.I. Vainshtein and V.I. Zakharov: *Nucl. Phys. B* **147**, 385 (1979); B.V. Geshkenbein, B.L. Ioffe and K.N. Zyblyuk: *Phys. Rev. D* **64**, 093009 (2001)
- [10] A.C. Kalloniatis, D.B. Leinweber, W. Melnitchouk and A.G. Williams, *Lattice Hadron Physics*. Proceedings, (Cairns, Australia, July 9-18, 2001)
- [11] C. Bernard, A. Duncan, J. LoSecco and S. Weinberg: *Phys. Rev. D* **12**, 792 (1975); E. Poggio, H. Quinn and S. Weinberg: *Phys. Rev. D* **13**, 1958 (1976); R. Shankar: *Phys. Rev. D* **15**, 755 (1977); K.G. Chetyrkin, N.V. Krasnikov and A.N. Tavkhelidze: *Phys. Lett. B* **76**, 83 (1978)
- [12] N.V. Krasnikov, A.A. Pivovarov and N. Tavkhelidze: *JETP Lett.* **36**, 333 (1982), *Z. Phys. C* **19**, 301 (1983); N.V. Krasnikov and A.A. Pivovarov: *Phys. Lett. B* **112**, 397 (1982); A.A. Pivovarov: *Phys. Atom. Nucl.* **62**, 1924 (1999); A.A. Pivovarov and A.S. Zubov: *Phys. Atom. Nucl.* **63**, 1650 (2000)
- [13] K. Schilcher and M.D. Tran: *Phys. Rev. D* **29**, 570 (1984); E. Braaten: *Phys. Rev. Lett.* **60**, 1606 (1988); S. Narison and A. Pich: *Phys. Lett. B* **211**, 183 (1988); A.A. Pivovarov: *Sov. J. Nucl. Phys.* **54**, 676 (1991); E. Braaten, S. Narison and A. Pich: *Nucl. Phys. B* **373**, 581 (1992)
- [14] S.G. Gorishny, A.L. Kataev and S.A. Larin: *Phys. Lett. B* **259**, 144 (1991); L.R. Surguladze and M.A. Samuel: *Phys. Rev. Lett.* **66**, 560 (1991); *Phys. Rev. D* **44**, 1602 (1991); K.G. Chetyrkin: *Phys. Lett. B* **391**, 402 (1997)
- [15] P.A. Baikov, K.G. Chetyrkin and J.H. Kühn: *Phys. Rev. Lett.* **88**, 012001 (2002)
- [16] K.G. Chetyrkin and A.A. Pivovarov: *Nuovo Cim. A* **100**, 899 (1988); A.A. Pivovarov: *JETP Lett.* **55**, 6 (1992); *Int. J. Mod. Phys. A* **10**, 3125 (1995)
- [17] ALEPH collaboration: *Z. Phys. C* **76**, 15 (1997); *Eur. Phys. J. C* **4**, 409 (1998); *ibid C* **11**, 599 (1999); OPAL collaboration: *Eur. Phys. J. C* **7**, 571 (1999)
- [18] J.G. Körner, F. Krajewski and A.A. Pivovarov: *Phys. Rev. D* **63**, 036001 (2001)

- [19] S. Groote, J.G. Körner, A.A. Pivovarov and K. Schilcher: Phys. Rev. Lett. **79**, 2763 (1997); S.J. Brodsky, J.R. Pelaez and N. Toumbas: Phys. Rev. D **60**, 037501 (1999); J.G. Körner, F. Krajewski and A.A. Pivovarov: Eur. Phys. J. C **12**, 461 (2000); *ibid* C **14**, 123 (2000)
- [20] K.G. Chetyrkin, J.H. Kühn and A.A. Pivovarov: Nucl. Phys. B **533**, 473 (1998); A. Pich and J. Prades: JHEP **9806**, 013 (1998); *ibid* **9910**, 004 (1999)
- [21] S. Narison and E. de Rafael: Phys. Lett. B **103**, 57 (1981); J. Gasser and H. Leutwyler: Phys. Rept. **87**, 77 (1982)
- [22] A.L. Kataev, N.V. Krasnikov and A.A. Pivovarov: Phys. Lett. B **123**, 93 (1983); Nuovo Cim. A **76**, 723 (1983)
- [23] J.G. Körner, F. Krajewski and A.A. Pivovarov: Eur. Phys. J. C **20**, 259 (2001)
- [24] A.L. Kataev and V.V. Starshenko: Mod. Phys. Lett. A **10**, 235 (1995); K.G. Chetyrkin, B.A. Kniehl and A. Sirlin: Phys. Lett. B **402**, 359 (1997)
- [25] A.A. Pivovarov: Z. Phys. C **53**, 461 (1992); Nuovo Cim. A **105**, 813 (1992); F. Le Diberder and A. Pich: Phys. Lett. B **286**, 147 (1992); S. Groote, J.G. Körner and A.A. Pivovarov: Phys. Lett. B **407**, 66 (1997); Mod. Phys. Lett. A **13**, 637 (1998)
- [26] G. Grunberg: Phys. Lett. B **95**, 70 (1980)
- [27] N.V. Krasnikov: Nucl. Phys. B **192**, 497 (1981); A.L. Kataev, N.V. Krasnikov and A.A. Pivovarov: Phys. Lett. B **107**, 115 (1981); Nucl. Phys. B **198**, 508 (1982); A. Dhar and V. Gupta: Phys. Rev. D **29**, 2822 (1984)
- [28] W.J. Marciano and A. Sirlin: Phys. Rev. Lett. **61**, 1815 (1988); E. Braaten and C.S. Li: Phys. Rev. D **42**, 3888 (1990)
- [29] C. Becchi, S. Narison, E. de Rafael and F.J. Yndurain: Z. Phys. C **8**, 335 (1981); A.A. Ovchinnikov and A.A. Pivovarov: Phys. Lett. B **163**, 231 (1985); N.V. Krasnikov and A.A. Pivovarov: Nuovo Cim. A **81**, 680 (1984); Y. Chung et al.: Z. Phys. C **25**, 151 (1984); H.G. Dosch and S. Narison: Phys. Lett. B **417**, 173 (1998)
- [30] A.A. Pivovarov: “Spectrality, coupling constant analyticity and the renormalization group,” to appear in Yad. Fiz. [arXiv:hep-ph/0104213]; S. Groote, J.G. Körner and A.A. Pivovarov: Phys. Rev. D **65**, 036001 (2002)
- [31] S.G. Gorishnii, A.L. Kataev and S.A. Larin: Phys. Lett. B **135**, 457 (1984); M. Jamin and M. Munz: Z. Phys. C **66**, 633 (1995); K.G. Chetyrkin, D. Pirjol and K. Schilcher: Phys. Lett. B **404**, 337 (1997)
- [32] K. Maltman and J. Kambor: Phys. Rev. D **65**, 074013 (2002); Phys. Lett. B **517**, 332 (2001); Nucl. Phys. A **684**, 348 (2001); Phys. Rev. D **62**, 093023 (2000); K. Maltman: Nucl. Phys. Proc. Suppl. A **109**, 178 (2002); arXiv:hep-ph/0209091
- [33] S. Chen et al.: Eur. Phys. J. C **22**, 31 (2001)
- [34] ALPHA and UKQCD Collaboration (J. Garden et al.): Nucl. Phys. B **571**, 237 (2000); ALPHA Collaboration (M. Guagnelli et al.): Nucl. Phys. B **560**, 465 (1999)